

Quiz - may 2024

1. MCMA twinning occurs if division occurs

- A) before 24 hrs
- B) 1-4 days
- C) 4-8 days
- D) >8 days.

Ans: D

2. what is the most important cause of perinatal mortality in twins?

- A) prematurity
- B) FGR
- C) TTTTS
- D) single fetal demise.

Ans: A

3. drug used in STOPPIT trial is

- A) progesterone
- B) steroids
- C) mgso4
- D) oxytocin.

Ans : B

4. according to hellin's law, the chances of twins in pregnancy are

- A) 1 in 60
- B) 1 in 70
- C) 1 in 80
- D) 1 in 100.

Ans : C

5. Important predictor of neurological outcome of the survivor after death of co twin is

- A) chorionicity
- B) gestational age at time of demise
- C) malformations present in deceased twin
- D) time between demise and delivery

Ans : A

6. the most common fetal congenital anomaly is

- A) down syndrome
- B) cardiac defect
- C) spina bifida
- D) cleft lip/ palate.

Ans : B

7. in twin reversed arterial perfusion donor is at risk of cardiomegaly and high output heart failure

- A) true.
- B) false

Ans : A

8. for twin discordance, the differences in the two twins should be 25% with the smaller twin as index.

A) true.                      B) false

Ans : B

9. fetal prognosis depends more on chorionicity and not zygosity

A) true.                      B) false

Ans : A

10. monozygotic twins are always monochorionic

A) true.                      B) false

Ans : B