## Mass per abdomen

<ol> <li>Which of the following a) unilocular cyst.</li> <li>thick septations.</li> </ol>	b) absent stror	mal invasion	ggests a malignant c	ovarian mass ??
2.Tumor marker co a) CA 19-9. c) CEA.	ommonly elevated b) CA 125 d) AFP.	-	/arian cancer?	
<ul><li>3. A woman with hadiagnosis?</li><li>a) leiomyoma.</li><li>c) functional ov cys</li></ul>	b) metasta	asis	vic mass.which of the	e following is most likely
<ul><li>4.A solid and cystic</li><li>a) observation.</li><li>c) CA 125.</li></ul>		сору	s lady. Next step in r	nanagement?
<ul><li>5. Characteristic U</li><li>a)Ground glass ap</li><li>c) honeycomb app</li></ul>	pearance b)tip of	iceberg sign		
<ul><li>6. Young woman w</li><li>a) serous cystader</li><li>b) yolk sac tumor.</li></ul>	noma. c) endom	netrioma	FP likely diagnosis?	
<ul><li>7. MC cause of abo</li><li>a) ovarian cancer.</li><li>c) TO abscess.</li></ul>		ibroid	ged woman?	
<ul><li>8. PLCO trial which</li><li>a) prostate.</li><li>c) ovary.</li></ul>	n cancer is not ind b) lung. d) cervix.	cluded? Ans : d		
<ul><li>9. Which is the dos</li><li>a) nephrotoxicity.</li><li>c) neuropathy.</li></ul>	be limiting toxicity b) myelosupp d.)ototoxicity	pression	he treatment of ovar	an cancer?

10. Newer drug which shrink fibroid by inhibiting growth factor and angiogenesis?

a) Elagolix. b) ulipristal acetate

c) mifepristone. d) ormeloxifene

Ans :a